

Production and β half lives of heavy neutron-rich nuclei approaching the r-process path at N=126



José Benlliure



Universidad de Santiago de Compostela
Spain

Nuclear data for the r process

Ground state properties of nuclei involved in the r process such as β half lives or masses are required for the full understanding of this nucleosynthesis mechanism.

Present RIB facilities made possible to produce light and medium-mass neutron-rich nuclei at the r-process path. However, the region around the A~195 waiting point is out of our reach.

The waiting point at A~195 defines the abundance of the heaviest elements in the Universe. But this is also an interesting region for nuclear structure because of the interplay between shell closure and deformation effects.



Nucler data for the r process



The β half lives of r-process nuclei along the N=126 shell define the role of the A~195 waiting point in the r process:

- ✓ Matter flow through the N=126 bottleneck region fixing the abundance pattern of the heaviest elements in the Universe.
- ✓ The velocity of synthesis of these heavy elements: r-process end point, r-process cycling.

Present theoretical predictions of the β half lives of r process nuclei close or at N=126 are rather discordant.

It is our goal to investigate:

- the production of heavy neutron-rich nuclei
- determine their β half lives



[1] P. Möller, et al. PRC 67, 055802 (2003)

[2] I. N. Borzov PRC 67, 025802 (2003)

energies

Target-spectator

Target


Fireball

Projectile-spectator

Projectile

β_0

β_0





- neutron-deficient nuclei are highly populated up to the drip-line
- the in-flight fragmentation of heavy nuclei leading to fission produces medium-mass neutron-rich nuclei
- the large fluctuation in isospin (abrasion) and excitation energy (ablation) give access to **cold fragmentation** processes where neutron-rich nuclei are produced.

Production of heavy neutron-rich nuclei

Experimental technique: SIS18+FRS (GSI)


^{238}U , $^{208}\text{Pb}(1 \text{ A GeV}) + \text{Be}$ (10^7 ions/s)




✓ One of the challenges of the experiment was the identification of charge states

→ Beam energy ($< 700 \text{ A MeV}$)

→ FRS+energy degrader






Production of heavy neutron-rich nuclei





75 heavy neutron-rich nuclei have been identified for the first time.

Production of heavy neutron-rich nuclei




β half lives


Experimental technique: ion – β ($-\gamma$) correlations



Active stopper: highly segmented DSSSD
ion- β time-position correlation



RISING γ -ray spectrometer
 β - γ prompt correlation





β half lives

Time correlations evaluation

Standard exponential fits of time correlations not applicable


- ✓ pulsed beam: time-modulated δ -electron background
- ✓ half lives (~ 10 s) are longer than the production/implantation rate: multiple implantations



New proposed method:


- ✓ β - γ prompt correlations: δ -electron suppression
- ✓ backward-time ion- β correlations: remaining background evaluation
- ✓ numerical fitting function based on Monte Carlo simulations of the implantation-decay process including experimental implantation rates and having as free parameters the β decay half life and the β detection efficiency

T. Kurtukian-Nieto et al., NIMA 67, 055802 (2008)




β half lives

Time correlations evaluation



β half lives

Results



The β half lives of 13 heavy neutron-rich nuclei have been determined, 11 of them for the first time.


Nuclei	w/o γ	with γ	other works	FRDM+ QRPA ^[1]	DF3+ QRPA ^[2]
²⁰⁴ Au		37 ± 0.8 s	39.8 ± 0.9 s		
²⁰⁴ Pt		16^{+6}_{-5} s		321.8 s	7.4 s
²⁰³ Pt		22 ± 4 s		654.0 s	12.7 s
²⁰² Ir	11 ± 3 s	15 ± 3 s		68.4 s	9.8 s
²⁰¹ Ir		21 ± 5 s		130.0 s	28.4 s
²⁰⁰ Ir		43^{+6}_{-5} s		124.1 s	25.0 s
¹⁹⁹ Ir	6^{+5}_{-4} s			370.6 s	46.7 s
¹⁹⁸ Ir	8 ± 2 s		8 ± 1 s	377.1 s	19.1 s
²⁰⁰ Os	6^{+4}_{-3} s			187.1 s	6.9 s
¹⁹⁹ Os	5^{+4}_{-2} s			106.8 s	6.6 s
¹⁹⁶ Re	3^{+1}_{-2} s			3.6 s	1.4 s
¹⁹⁵ Re	6 ± 1 s			3.3 s	8.5 s
¹⁹⁴ Re	1 ± 0.5 s			70.8 s	2.1 s

[1] P. Möller, et al. PRC 67, 055802 (2003)


[2] I. N. Borzov PRC 67, 025802 (2003)

β half lives

Results



- ✓ FRDM calculations overestimate the measured half lives
- ✓ DF3 calculations provides a better description of the data: role of first forbidden transitions
- ✓ These results may suggest that the r process matter flow to the heavier fissioning nuclei could be faster than expected



Summary

- The production of heavy neutron-rich nuclei close to the $A \sim 195$ r process waiting point was investigated using fragmentation reactions of ^{238}U and ^{208}Pb beams at relativistic energies.
 - ✓ 75 neutron-rich isotopes of elements between Yb and Fr were identified for the first time
 - ✓ Their production cross sections were determined and used to benchmark model calculations

- β half lives were determined using ion – β ($-\gamma$) correlations
 - ✓ A new method to determine β half lives under complex background conditions was introduced
 - ✓ The half lives of 13 heavy neutron-rich nuclei were determined.
 - ✓ The measured half lives are overestimated by FRDM+QRPA calculations and rather well described by DF3+QRPA.
 - ✓ The confirmation of this result for r process nuclei would indicate a faster r process matter flow towards the heavier fissioning nuclei.

Collaborators

J. Benlliure¹, **A.I. Morales¹**, **T. Kurtukian-Nieto^{1,a}**, **H. Alvarez-Pol¹**, I.N. Borzov²,
 L. Audouin³, B. Blank⁴, F. Becker², E. Casarejos¹, D. Cortina-Gil¹, T. Enqvist^{2,b},
 B. Fernández⁵, M. Fernández-Ordóñez^{1,c}, J. Giovinazzo⁴, D. Henzlova^{2,d}, A.R. Junghans⁶,
 B. Jurado⁴, k. Langanke², G. Martinez-Pinedo², P. Napolitani³, J. Pereira^{1,d}, F. Rejmund⁵,
 K.-H. Schmidt², O. Yordanov^{2,e}

¹ Departamento de Física de Partículas, Universidade de Santiago de Compostela

² GSI Darmstadt

³ Institut de Physique Nucleaire d'Orsay

⁴ Centre d'Etudes Nucleaires Bordeaux-Gradignan

⁵ GANIL, Caen

⁶ Forschungszentrum Dresden-Rossendorf

and the RISING collaboration